

Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

2.4.2 Population size (other than individuals)	Unit	N/A
	min	max
2.4.3 Additional information	Definition of locality	
	Conversion method	
	Problems	
2.4.4 Year or period	2012	
2.4.5 Method – population size	Complete survey/Complete survey or a statistically robust estimate (3)	
2.4.6 Short-term trend period	2001-2012	
2.4.7 Short term trend direction	increase (+)	
2.4.8 Short-term trend magnitude	min	max confidence interval
2.4.9 Short-term trend method	Estimate based on expert opinion with no or minimal sampling (1)	
2.4.10 Long-term trend period		
2.4.11 Long term trend direction	N/A	
2.4.12 Long-term trend magnitude	min	max confidence interval
2.4.13 Long-term trend method	N/A	
2.4.14 Favourable reference population	number	800
	operator	N/A
	unknown	No
	method	The favourable reference population is defined as approximately the minimum number of individuals counted in 2014 (rounded to the upper number of individuals).
2.4.15 Reason for change	Improved knowledge/more accurate data	

2.5 Habitat for the Species

2.5.1 Surface area - Habitat (km ²)	0,06
2.5.2 Year or period	2012
2.5.3 Method used - habitat	Complete survey/Complete survey or a statistically robust estimate (3)
2.5.4 a) Quality of habitat	Good
2.5.4 b) Quality of habitat - method	Clinopodium taygeteum grows on hard or eroded limestone rocks and cliffs. There are several rock or cliff systems in the distribution range of the species, some of them partially or entirely inaccessible. This habitat is of good quality regarding species composition (typical species) and because it is not significantly influenced by grazing.
2.5.5 Short term trend period	2001-2012
2.5.6 Short term trend direction	unknown (x)
2.5.7 Long-term trend period	
2.5.8 Long term trend direction	N/A
2.5.9 Area of suitable habitat (km ²)	0
2.5.10 Reason for change	

2.6 Main Pressures

Pressure	ranking	pollution qualifier(s)
non intensive grazing (A04.02)	low importance (L)	N/A
burning down (J01.01)	low importance (L)	N/A

2.6.1 Method used – pressures based exclusively or to a larger extent on real data from sites/occurrences or other

2.7 Main Threats

Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

Threat	ranking	pollution qualifier(s)
non intensive grazing (A04.02)	low importance (L)	N/A
burning down (J01.01)	low importance (L)	N/A
mountaineering & rock climbing (G01.04.01)	low importance (L)	N/A

2.7.1 Method used – threats expert opinion (1)

2.8 Complementary Information

2.8.1 Justification of % thresholds for trends

2.8.2 Other relevant Information

Note on 2.4.2. Population size was estimated in 2014 and c. 760 individuals were counted. This value is reported as minimum population size. The species often grows at inaccessible rocks and it is certain that the population size is larger. Since the species was observed to have an uneven occurrence pattern in its distribution area, it is difficult to predict the maximum number of individuals. Taking into consideration the habitat of the species, not rare on the mountain, a prediction of c. 2.000 mature individuals can serve as a maximum population size. Note on 2.4.15. According to new data, the population size of *Clinopodium taygeteum* is larger than the size reported in 2006. This difference is purely due to a better estimation of population size in the field (several localities visited, more time allocated to estimations in situ) and should not be attributed to a genuine increase in the number of individuals. The total population is presumably stable over a long time.

Note on 2.5.1: Habitat area estimation is the sum of the area actually covered by the population of the species.

The presence of the species in GR2520005, GR2520006 is not reported in the literature or by any other source. Possible discrepancies with the SDF will be corrected in the next submission of revised SDF forms

2.8.3 Trans-boundary assessment

2.9 Conclusions (assessment of conservation status at end of reporting period)

2.9.1 Range assessment Favourable (FV)
qualifiers N/A

2.9.2. Population assessment Favourable (FV)
qualifiers N/A

2.9.3. Habitat assessment Favourable (FV)
qualifiers N/A

2.9.4. Future prospects assessment Favourable (FV)
qualifiers N/A

2.9.5 Overall assessment of Conservation Status Favourable (FV)

2.9.5 Overall trend in Conservation Status N/A

3. Natura 2000 coverage and conservation measures - Annex II species

3.1 Population

3.1.1 Population Size

Unit	number of individuals (i)
min	80
max	100

Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B)

3.1.2 Method used

Complete survey/Complete survey or a statistically robust estimate (3)

3.1.3 Trend of population size within

unknown (x)

3.2 Conservation Measures

3.2.1 Measure	3.2.2 Type	3.2.3 Ranking	3.2.4 Location	3.2.5 Broad Evaluation
Legal protection of habitats and species (6.3)	Legal	high importance (H)	Inside	Maintain Long term
Regulation/ Management of hunting and taking (7.1)	Legal	high importance (H)	Inside	Long term