

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

CODE: 5110

NAME: Stable xerothermophilous formations with *Buxus sempervirens* on rock slopes (Berberidion p.p.)

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.2 Distribution Method	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	2006-2012
1.1.4 Additional map	No
1.1.5 Range Map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published

Mediterranean (MED)

Dimopoulos P., Xystrakis F. and Tsiripidis I. 2014. Deliverable A1. Final Catalogue of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 54.

Dimopoulos P., Fotiadis G., Tsiripidis I., Panitsa M. and Karadimou E. 2014. Deliverable A2. Report and Literature Database on Habitat Types of Greece – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 210.

Tsiripidis I., Xystrakis F., Kasampalis D., Mastrogianni A., Strid A. and Dimopoulos P., 2014. Deliverable A4. Potential Distribution Maps of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, Athens, pages 176.

Dimopoulos P., Tsiripidis I., Xystrakis F., Panitsa M., Fotiadis G., Kallimanis A.S. and Kazoglou I. 2014. Deliverable A6. Explanatory Implementation Manual for the Conservation Degree Assessment of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 35. (with Annexes: I. Habitat types protocols, pages 600; II. Explanatory notes on the habitat types protocols selection, pages 4; III. Correspondence of Habitat types protocols with the clusters of vegetation relevés (excel file).

Dimopoulos P., Tsiripidis I., Xystrakis F., Kallimanis A.S and Panitsa M. 2014. Deliverable A7. Preliminary Analysis of the Field Data for the Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 16.

2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km ²)	300
2.3.2 Range method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.3.3 Short-term trend period	2001-2012
2.3.4 Short-term trend direction	stable (0)
2.3.5 Short-term trend magnitude	min max
2.3.6 Long-term trend period	
2.3.7 Long-term trend direction	N/A
2.3.8 Long-term trend magnitude	min max
2.3.9 Favourable reference range	area (km ²) operator approximately equal to (≈) unkown No method
2.3.10 Reason for change	Improved knowledge/more accurate data Use of different method

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.4 Area covered by Habitat

2.4.1 Surface area (km ²)	14,13
2.4.2 Year or period	2000-2012
2.4.3 Method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.4.4 Short-term trend period	2001-2012
2.4.5 Short-term trend direction	stable (0)
2.4.6 Short-term trend magnitude	min max
2.4.7 Short term trend method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.4.8 Long-term trend period	
2.4.9 Long-term trend direction	N/A
2.4.10 Long-term trend magnitude	min max
2.4.11 Long term trend method used	N/A
2.4.12 Favourable reference area	area (km) operator approximately equal to (≈) unknown No method
2.4.13 Reason for change	Improved knowledge/more accurate data Use of different method

2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
grazing (A04)	low importance (L)	N/A
Structures, buildings in the landscape (E04)	low importance (L)	N/A
fire and fire suppression (J01)	low importance (L)	N/A

2.5.1 Method used – pressures mainly based on expert judgement and other data (2)

2.6 Main Threats

Threat	ranking	pollution qualifier(s)
grazing (A04)	low importance (L)	N/A
Structures, buildings in the landscape (E04)	low importance (L)	N/A
fire and fire suppression (J01)	low importance (L)	N/A

2.6.1 Method used – threats expert opinion (1)

2.7 Complementary Information

2.7.1 Species

Acer hyrcanum

Allium heldreichii

Amelanchier ovalis

Aubrieta thessala

Berberis cretica

Buxus sempervirens

Carex distachya

Carex halleriana

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Cotoneaster nebrodensis

Cyclamen hederifolium

Daphne oleoides

Euonymus verrucosus

Fragaria viridis

Fritillaria messanensis

Galium hellenicum

Hippocrepis emerus

Juniperus communis

Juniperus oxycedrus

Lilium chalcedonicum

Lonicera etrusca

Orchis quadripunctata

Ostrya carpinifolia

Polygonatum odoratum

Pteridium aquilinum

Rhamnus saxatilis

Saxifraga chrysospleniifolia

Silene italica

Sorbus aria

Teucrium chamaedrys

Thymus sibthorpii

2.7.2 Species method used

Typical species were determined on the basis of a vegetation database, comprised of about 22000 sampling plots. First, a list of possible typical species was determined per habitat type, selecting the ones presenting a high fidelity value to the habitat types according the algorithm of Tsiripidis et al. (2009) and the phi coefficient value (Chytrý et al. 2002). Then typical species per habitat type were selected from the above-mentioned lists by expert judgment and using as criteria species niche breadth, their ability to comprise indicators of habitat types' conservation status and their function as keystone species. The nomenclature of the typical species follows Dimopoulos et al. (2013).ReferencesChytrý, M., Tichý, L., Holt, J. & Botta-Dukát, J. 2002. Determination of diagnostic species with statistical fidelity measures. *Journal of Vegetation Science* 13: 79–90. Dimopoulos, P., Raus, Th., Bergmeier, E., Constantinidis, Th., Iatrou, G., Kokkini, S., Strid, A. & Tzanoudakis, D. 2013: *Vascular plants of Greece: an annotated checklist*. – Berlin: Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin; Athens: Hellenic Botanical Society. *Englera* 31: 1-367. Tsiripidis, I., Bergmeier, E., Fotiadis, G. & Dimopoulos, P. 2009. A new algorithm for the determination of differential taxa. *Journal of Vegetation Science* 20: 233-240.

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.7.3 Justification of % - thresholds for trends

2.7.4 Structure and functions - methods used

Complete survey/Complete survey or a statistically robust estimate (3)

2.7.5 Other relevant information

2.8 Conclusions (assessment of conservation status at end of reporting period)

2.8.1 Range

assessment Favourable (FV)
qualifiers N/A

2.8.2 Area

assessment Favourable (FV)
qualifiers N/A

2.8.3 Specific structures and functions (incl Species)

assessment Favourable (FV)
qualifiers N/A

2.8.4 Future prospects

assessment Favourable (FV)
qualifiers N/A

2.8.5 Overall assessment of Conservation Status

Favourable (FV)

2.8.5 Overall trend in Conservation Status

N/A

3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

3.1 Area covered by habitat

3.1.1 Surface area (km²)

min 4,6 max 4,6

3.1.2 Method used

Complete survey/Complete survey or a statistically robust estimate (3)

3.1.3. Trend of surface area

stable (0)

3.2 Conversation Measures

3.2.1 Measure

3.2.2 Type

3.2.3 Ranking

3.2.4 Location

3.2.5 Broad Evaluation

Establish protected areas/sites (6.1)

Legal
Administrative
One-off

medium
importance (M)

Inside

Maintain
Long term