

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

CODE: 9260

NAME: Castanea sativa woods

1. National Level

1.1 Maps

1.1.1 Distribution Map	Yes
1.1.2 Distribution Method	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	2006-2012
1.1.4 Additional map	No
1.1.5 Range Map	Yes

2. Biogeographical Or Marine Level

2.1 Biogeographical Region

2.2 Published

Mediterranean (MED)

Dimopoulos P., Xystrakis F. and Tsiripidis I. 2014. Deliverable A1. Final Catalogue of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 54.

Dimopoulos P., Fotiadis G., Tsiripidis I., Panitsa M. and Karadimou E. 2014. Deliverable A2. Report and Literature Database on Habitat Types of Greece – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 210.

Tsiripidis I., Xystrakis F., Kasampalis D., Mastrogianni A., Strid A. and Dimopoulos P., 2014. Deliverable A4. Potential Distribution Maps of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, Athens, pages 176.

Dimopoulos P., Tsiripidis I., Xystrakis F., Panitsa M., Fotiadis G., Kallimanis A.S. and Kazoglou I. 2014. Deliverable A6. Explanatory Implementation Manual for the Conservation Degree Assessment of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 35. (with Annexes: I. Habitat types protocols, pages 600; II. Explanatory notes on the habitat types protocols selection, pages 4; III. Correspondence of Habitat types protocols with the clusters of vegetation relevés (excel file).

Dimopoulos P., Tsiripidis I., Xystrakis F., Kallimanis A.S and Panitsa M. 2014. Deliverable A7. Preliminary Analysis of the Field Data for the Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 16.

Barbero M. & Quézel P. 1976. Les groupements forestiers de Grece Centro-Meridionale. *Ecologia Mediterranea* 2: 1-86.

Bergmeier E. 1990. Walder und Gebusche des Niederen Olymp (Kato Olimbos, NO-Thessalien). *Phytocoenologia* 18(2/3): 161-342.

Δημητράλλος Ν.Γ. 2005. Γεωβοτανική Έρευνα του Όρους Τυμφρηστού (ΒΔ Στερεά Ελλάδα) Χλωρίδα - Βλάστηση - Αξιολόγηση - Διαχείριση. Διδακτορική Διατριβή. Πανεπιστήμιο Πατρών, σελ. 296.

Dimopoulos P., Bergmeier E., Theodoropoulos K. & Eleftheriadou E. 2005. Thermophilous deciduous forests in Greece – a preliminary survey. *Bot. Chron.* 18(1): 83-100.

Gamisans J. & Hebrard J.-P. 1980. A propos de la vegetation des forets en Grece du Nord-est (Macedoine orientale et Thrace occidentale). *Documents phytosociologiques* 5:243-276

Gamisans J. & Hebrard J.-P. 1979. A propos de la vegetation des forets d' Epire et de Macedoine Grecque occidentale. *Documents phytosociologiques* IV: 290-327.

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

- Grandstein S.R. & Smittenberg J.H. 1977. The hydrophilus vegetation of western Crete. *Vegetatio* 34(2): 65-86.
- Θεοδωρόπουλος Κ. 2001. Ζώνες βλάστησης και τύποι οικοτόπων του νομού Θεσσαλονίκης. *Επιστ. Επετ. Τμημ. Δασολογίας & Φυσικού Περιβάλλοντος ΜΔ*: 353-381.
- Καραγιαννακίδου-Ιατροπούλου Β. 1983. Σταθμολογική έρευνα στην Quercus – Fagetea κλάση του ορεινού συγκροτήματος του Χορτιάτη. *Διδακτορική Διατριβή*. Α.Π.Θ. *Επιστ. Επετ. Σχολής Θετικών Επιστημών* 22(25), σελ. 161.
- Konstantinidis P., Tsiogrlis G., Xofis P. & Buckley G.P. 2008. Taxonomy and ecology of *Castanea sativa* Mill. *Forests in Greece*. *Plant Ecol.* 195: 235-256
- Κοράκης Γ.Χ. 2003. Οι μονάδες βλάστησης του όρους Πάικου και η αξιολόγησή τους από αναδασωτική σκοπιά. *Διδακτορική Διατριβή*. ΑΠΘ, σελ. 264 + Παράρτημα α.
- Κοράκης Γ. & Αθανασιάδης Ν. 2003. Οι φυτοκοινωνίες των δασών της *Castanea sativa* Miller στο όρος Πάικο. *Πρακτικά 11ου Πανελληνίου Δασολογικού Συνεδρίου της Ελληνικής Δασολογικής Εταιρείας, Αρχαία Ολυμπία, 30/9-3/10 2003*: 615-624.
- Κωνσταντινίδης Π. & Τσιουρλής Γ. 2001. Οι βλαστητικές μονάδες (τύποι οικοτόπων) της Επαρχίας Λαγκαδά (Λεκάνη Μυγδονίας): Μέρος Ι: Περιγραφή, ανάλυση και χαρτογράφηση. *Επιστ. Επετ. Τμημ. Δασολογίας & Φυσικού Περιβάλλοντος ΜΔ*: 627-654.
- Κωνσταντινίδης Π. & Τσιουρλής Γ. 2001. Οι τύποι οικοτόπων της Επαρχίας Λαγκαδά (Λεκάνης Μυγδονίας): Μέρος ΙΙ. Οικολογική κατάσταση και δυναμική. *Επιστ. Επετ. Τμημ. Δασολογίας & Φυσικού Περιβάλλοντος ΜΔ*: 655-680.
- Μαρούλης Γ. 2003. Χλωρίδα και βλάστηση των οικοσυστημάτων του όρους Ερύμανθος (ΒΔ Πελοπόννησος). *Διδακτορική Διατριβή*. Πανεπιστήμιο Πατρών, σελ. 450 + 1 Παράρτημα + 1 Χάρτης.
- Μπαζός Ι. & Γιαννίτσαρος Α. 2005. Χλωρίδα και βλάστηση της Λέσβου: γενική επισκόπηση. *Πρακτικά 10ου Πανελληνίου Επιστημονικού Συνεδρίου της Ελληνικής Βοτανικής Εταιρείας, Ιωάννινα, 5-8 Μαΐου 2005*, σελ. 8 (σε CD).
- Ντάφης Σ. 1966. Σταθμολογικά και δασοαποδοτικά έρευναι εις πρεμνοφυή δρυοδάση και καστανωτά της βορειοανατολικής Χαλκιδικής. *Θεσσαλονίκη*, σελ. 113.
- Raus Th. 1980. Die vegetation Osthessaliens (Griechenland), III. Quercus-Fagetea und azonale Gehölzgesellschaften. *Bot. Jahrb. Syst.* 101(3): 313-361.
- Φωτιάδης Γ. 2004. Καθορισμός των δασικών φυτοκοινωνιολογικών μονάδων του Ελληνικού τμήματος του όρους Μπέλες και της οροσειράς των Κρουσίων. *Διδακτορική Διατριβή*, ΑΠΘ, σελ. 273 + Παράρτημα.
- Zoller H., Geissler P. & Athanasiadis N. 1977. Beiträge zur Kenntnis der Wälder, Moos- und Flechtenassoziationen in den Gebirgen Nordgriechenlands. *Bauhinia* 6/1: 215-255.

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km ²)	2229
2.3.2 Range method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.3.3 Short-term trend period	2001-2012
2.3.4 Short-term trend direction	stable (0)
2.3.5 Short-term trend magnitude	min max
2.3.6 Long-term trend period	
2.3.7 Long-term trend direction	N/A
2.3.8 Long-term trend magnitude	min max
2.3.9 Favourable reference range	area (km ²) operator approximately equal to (≈) unkown No method
2.3.10 Reason for change	Improved knowledge/more accurate data Use of different method

2.4 Area covered by Habitat

2.4.1 Surface area (km ²)	484,2
2.4.2 Year or period	2000-2012
2.4.3 Method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.4.4 Short-term trend period	2001-2012
2.4.5 Short-term trend direction	stable (0)
2.4.6 Short-term trend magnitude	min max
2.4.7 Short term trend method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.4.8 Long-term trend period	
2.4.9 Long-term trend direction	N/A
2.4.10 Long-term trend magnitude	min max
2.4.11 Long term trend method used	N/A
2.4.12 Favourable reference area	area (km) operator approximately equal to (≈) unknown No method
2.4.13 Reason for change	Improved knowledge/more accurate data Use of different method

2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
modification of cultivation practices (A02)	low importance (L)	N/A
grazing (A04)	low importance (L)	N/A
Forest and Plantation management & use (B02)	low importance (L)	N/A
Roads, paths and railroads (D01)	low importance (L)	N/A
Outdoor sports and leisure activities, recreational activities (G01)	low importance (L)	N/A
invasive non-native species (I01)	low importance (L)	N/A
Interspecific faunal relations (K03)	low importance (L)	N/A

2.5.1 Method used – pressures mainly based on expert judgement and other data (2)

2.6 Main Threats

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Threat	ranking	pollution qualifier(s)
modification of cultivation practices (A02)	low importance (L)	N/A
grazing (A04)	low importance (L)	N/A
Forest and Plantation management & use (B02)	low importance (L)	N/A
invasive non-native species (I01)	low importance (L)	N/A
fire and fire suppression (J01)	low importance (L)	N/A
Interspecific faunal relations (K03)	low importance (L)	N/A

2.6.1 Method used – threats expert opinion (1)

2.7 Complementary Information

2.7.1 Species

Acer hyrcanum

Aremonia agrimonoides

Brachypodium sylvaticum

Calamintha grandiflora

Campanula spatulata

Castanea sativa

Clinopodium vulgare

Cornus mas

Cytisus villosus

Euphorbia amygdaloides

Fragaria vesca

Fraxinus ornus

Hedera helix

Lathyrus laxiflorus

Lathyrus niger

Luzula forsteri

Melica uniflora

Physospermum cornubiense

Poa nemoralis

Potentilla micrantha

Primula acaulis

Prunus avium

Pteridium aquilinum

Quercus frainetto

Sanicula europaea

Sorbus domestica

Sorbus torminalis

Tilia tomentosa

Verbascum nigrum

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Veronica chamaedrys

2.7.2 Species method used

Typical species were determined on the basis of a vegetation database, comprised of about 22000 sampling plots. First, a list of possible typical species was determined per habitat type, selecting the ones presenting a high fidelity value to the habitat types according the algorithm of Tsiripidis et al. (2009) and the phi coefficient value (Chytrý et al. 2002). Then typical species per habitat type were selected from the above-mentioned lists by expert judgment and using as criteria species niche breadth, their ability to comprise indicators of habitat types' conservation status and their function as keystone species. The nomenclature of the typical species follows Dimopoulos et al. (2013).

References

- Chytrý, M., Tichý, L., Holt, J. & Botta-Dukát, J. 2002. Determination of diagnostic species with statistical fidelity measures. *Journal of Vegetation Science* 13: 79–90.
- Dimopoulos, P., Raus, Th., Bergmeier, E., Constantinidis, Th., Iatrou, G., Kokkini, S., Strid, A. & Tzanoudakis, D. 2013: Vascular plants of Greece: an annotated checklist. – Berlin: Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin; Athens: Hellenic Botanical Society. *Englera* 31: 1-367.
- Tsiripidis, I., Bergmeier, E., Fotiadis, G. & Dimopoulos, P. 2009. A new algorithm for the determination of differential taxa. *Journal of Vegetation Science* 20: 233-240.

2.7.3 Justification of % - thresholds for trends

2.7.4 Structure and functions - methods used

2.7.5 Other relevant information

Complete survey/Complete survey or a statistically robust estimate (3)

2.8 Conclusions (assessment of conservation status at end of reporting period)

2.8.1 Range

assessment Favourable (FV)
qualifiers N/A

2.8.2 Area

assessment Favourable (FV)
qualifiers N/A

2.8.3 Specific structures and functions (incl Species)

assessment Favourable (FV)
qualifiers N/A

2.8.4 Future prospects

assessment Favourable (FV)
qualifiers N/A

2.8.5 Overall assessment of Conservation Status

Favourable (FV)

2.8.5 Overall trend in Conservation Status

N/A

3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

3.1 Area covered by habitat

3.1.1 Surface area (km²)

min 135,7 max 135,7

3.1.2 Method used

Complete survey/Complete survey or a statistically robust estimate (3)

3.1.3. Trend of surface area

stable (0)

Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

3.2 Conservation Measures

3.2.1 Measure	3.2.2 Type	3.2.3 Ranking	3.2.4 Location	3.2.5 Broad Evaluation
Establish protected areas/sites (6.1)	Legal Administrative One-off	medium importance (M)	Inside	Maintain Long term