

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

NATIONAL LEVEL

1. General information

1.1 Member State	GR
1.2 Habitat code	91F0 - Riparian mixed forests of <i>Quercus robur</i> , <i>Ulmus laevis</i> and <i>Ulmus mino</i>

2. Maps

2.1 Year or period	2015
2.3 Distribution map	Yes
2.3 Distribution map Method used	Based mainly on extrapolation from a limited amount of data
2.4 Additional maps	Yes

BIOGEOGRAPHICAL LEVEL

3. Biogeographical and marine regions

3.1 Biogeographical or marine region where the habitat occurs	Mediterranean (MED)
3.2 Sources of information	<p>Dimopoulos P., Xystrakis F. and Tsiropidis I. 2014. Deliverable A1. Final Catalogue of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 54.</p> <p>Dimopoulos P., Fotiadis G., Tsiropidis I., Panitsa M. and Karadimou E. 2014. Deliverable A2. Report and Literature Database on Habitat Types of Greece – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 210.</p> <p>Tsiropidis I., Xystrakis F., Kasampalis D., Mastrogianni A., Strid A. and Dimopoulos P., 2014. Deliverable A4. Potential Distribution Maps of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, Athens, pages 176.</p> <p>Dimopoulos P., Tsiropidis I., Xystrakis F., Panitsa M., Fotiadis G., Kallimanis A.S. and Kazoglou I. 2014. Deliverable A6. Explanatory Implementation Manual for the Conservation Degree Assessment of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 35. (with Annexes: I. Habitat types protocols, pages 600; II. Explanatory notes on the habitat types protocols selection, pages 4; III. Correspondence of Habitat types protocols with the clusters of vegetation relevés (excel file).</p> <p>Dimopoulos P., Tsiropidis I., Xystrakis F., Kallimanis A.S and Panitsa M. 2014. Deliverable A7. Preliminary Analysis of the Field Data for the Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 16.</p> <p>Αθανασιάδης Ν., Θεοδωρόπουλος Κ., Ελευθεριάδου Ε. & Δρόσος Ε. 1996. Δασικές φυτοκοινωνίες του δέλτα του Θεσσαλικού Πηνειού. Επιστ. Επετ. Τμημ. Δασολογίας και Φυσικού Περιβάλλοντος 39(2): 879-902.</p> <p>Αθανασιάδης Ν.Η. & Δρόσος Ε.Γ. 1989. <i>Leucujo-Fraxinetum parvifoliae</i> Glavac 59 και <i>Pruno-Fraxinetum</i> Oberd. 53 του Δέλτα του Θεσσαλικού Πηνειού. Επιστ. Επετ. Τμημ. Δασολογίας και Φυσικού Περιβάλλοντος 1Β(1): 543-558.</p> <p>Παναγιωτίδης Σ. & Φωτιάδης Γ. 2001. Μονάδες βλάστησης κατάλοιπων παρόχθιων δασών της κεντρικής Μακεδονίας. Επιστ. Επετ. Τμημ. Δασολογίας & Φυσικού Περιβάλλοντος ΜΔ: 477-488.</p> <p>Raus Th. 1980. Die vegetation Osthessaliens (Griechenland), III. <i>Querco-Fagetea</i> und azonale Gehölzgesellschaften. Bot. Jahrb. Syst. 101(3): 313-361.</p>

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

	b) Operator	Approximately equal to (\approx)
	c) Unknown	Yes
	d) Method	
5.14 Change and reason for change in surface area of range	No change	
	The change is mainly due to:	

5.15 Additional information

6. Structure and functions

6.1 Condition of habitat	a) Area in good condition (km ²)	Minimum 0	Maximum 0
	b) Area in not-good condition (km ²)	Minimum 28,92	Maximum 28,92
	c) Area where condition is not known (km ²)	Minimum 67,48	Maximum 67,48
6.2 Condition of habitat Method used	Complete survey or a statistically robust estimate		
6.3 Short-term trend of habitat area in good condition Period	20072018		
6.4 Short-term trend of habitat area in good condition Direction	Increasing (+)		
6.5 Short-term trend of habitat area in good condition Method used	Complete survey or a statistically robust estimate		
6.6 Typical species	Has the list of typical species changed in comparison to the previous reporting period? No		
6.7 Typical species Method used	<p>Typical species were determined on the basis of a vegetation database, comprised of about 22000 sampling plots. First, a list of possible typical species was determined per habitat type, selecting the ones presenting a high fidelity value to the habitat types according the algorithm of Tsiripidis et al. (2009) and the phi coefficient value (Chytrý et al. 2002). Then typical species per habitat type were selected from the above-mentioned lists by expert judgment and using as criteria species niche breadth, their ability to comprise indicators of habitat types' conservation status and their function as keystone species. The nomenclature of the typical species follows Dimopoulos et al. (2013). References Chytrý, M., Tichý, L., Holt, J. & Botta-Dukát, J. 2002. Determination of diagnostic species with statistical fidelity measures. <i>Journal of Vegetation Science</i> 13: 79–90. Dimopoulos, P., Raus, Th., Bergmeier, E., Constantinidis, Th., Iatrou, G., Kokkini, S., Strid, A. & Tzanoudakis, D. 2013: Vascular plants of Greece: an annotated checklist. – Berlin: Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin; Athens: Hellenic Botanical Society. Englera 31: 1-367. Tsiripidis, I., Bergmeier, E., Fotiadis, G. & Dimopoulos, P. 2009. A new algorithm for the determination of differential taxa. <i>Journal of Vegetation Science</i> 20: 233-240.</p>		
6.8 Additional information	Assumption: 20% of habitat area is estimated to be in not-good condition.		

7. Main pressures and threats

7.1 Characterisation of pressures/threats

Pressure	Ranking
Conversion into agricultural land (excluding drainage and burning) (A01)	H

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

Droughts and decreases in precipitation due to climate change (N02) M

Threat Ranking

Conversion into agricultural land (excluding drainage and burning) (A01) H

Drainage (K02) H

7.2 Sources of information PRESSURES: Based mainly on expert judgement and other data.
THREATS: Based on expert opinion.

7.3 Additional information

8. Conservation measures

8.1 Status of measures

a) Are measures needed?	Yes
b) Indicate the status of measures	Measures identified, but none yet taken

8.2 Main purpose of the measures taken

8.3 Location of the measures taken

8.4 Response to the measures

8.5 List of main conservation measures

Prevent conversion of natural and semi-natural habitats, and habitats of species into agricultural land (CA01)

Restoration of Annex I forest habitats (CB08)

Adopt climate change mitigation measures (CN01)

Implement climate change adaptation measures (CN02)

8.6 Additional information

9. Future prospects

9.1 Future prospects of parameters

a) Range	Good
b) Area	Poor
c) Structure and functions	Bad

9.2 Additional information

10. Conclusions

10.1. Range	Favourable (FV)
10.2. Area	Unfavourable - Inadequate (U1)
10.3. Specific structure and functions (incl. typical species)	Unfavourable - Bad (U2)
10.4. Future prospects	Unfavourable - Bad (U2)
10.5 Overall assessment of Conservation Status	Unfavourable - Bad (U2)
10.6 Overall trend in Conservation Status	Stable (=)

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

10.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

Improved knowledge/more accurate data

Use of different method

The change is mainly due to: Improved knowledge/more accurate data

b) Overall trend in conservation status

No change

The change is mainly due to:

10.8 Additional information

11. Natura 2000 (pSCIs, SCIs, SACs) coverage for Annex I habitat types

11.1 Surface area of the habitat type inside the pSCIs, SCIs and SACs network (in km² in biogeographical/marine region)

a) Minimum

b) Maximum

c) Best single value 2,86

11.2 Type of estimate

Minimum

11.3 Surface area of the habitat type inside the network Method used

Complete survey or a statistically robust estimate

11.4 Short-term trend of habitat area in good condition within the network Direction

Stable (0)

11.5 Short-term trend of habitat area in good condition within network Method used

Complete survey or a statistically robust estimate

11.6 Additional information

The change in 11.1 (in comparison to the previous report) is due to the updated mapping datasets on terrestrial habitat types within the Natura 2000 network (pSCIs, SCIs and SACs), based on the most recent national project (results became available in 2018). As this project did not include the extended areas of the Natura 2000 sites, nor the newly proposed SCIs, the surface area reported is the minimum.

12. Complementary information

12.1 Justification of % thresholds for trends

12.2 Other relevant information